

SPECIALTY GUIDELINE MANAGEMENT

PROLIA (denosumab)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indications

1. Treatment of postmenopausal women with osteoporosis at high risk for fracture
2. Treatment to increase bone mass in men with osteoporosis at high risk for fracture
3. Treatment to increase bone mass in men at high risk for fracture receiving androgen deprivation therapy (ADT) for non-metastatic prostate cancer
4. Treatment to increase bone mass in women at high risk for fracture receiving adjuvant aromatase inhibitor therapy for breast cancer
5. Treatment of men and women with glucocorticoid-induced osteoporosis at high risk for fracture

B. Compendial Uses

Prevention or treatment of osteoporosis during androgen deprivation therapy for patients with high fracture risk

All other indications are considered experimental/investigational and are not a covered benefit.

II. CRITERIA FOR INITIAL APPROVAL

A. **Osteoporosis in Postmenopausal Women**

Authorization of 12 months may be granted to postmenopausal female members when ANY of the following criteria are met:

1. Member has a history of fragility fractures
2. Member has a pre-treatment T-score of ≤ -2.5 OR member has osteopenia with a high pre-treatment FRAX fracture probability (See Appendix B) and meets ANY of the following criteria:
 - a. Member has indicators of higher fracture risk (e.g., advanced age, frailty, glucocorticoid use, very low T-scores, or increased fall risk)
 - b. Member has failed prior treatment with or is intolerant to previous injectable osteoporosis therapy (e.g., zoledronic acid [Reclast], teriparatide [Forteo])
 - c. Member has had an oral bisphosphonate trial of at least 1-year duration or there is a clinical reason to avoid treatment with an oral bisphosphonate (See Appendix A)

B. **Osteoporosis in Men**

Authorization of 12 months may be granted to male members with osteoporosis when ANY of the following criteria are met:

1. Member has a history of an osteoporotic vertebral or hip fracture
2. Member has a pre-treatment T-score of ≤ -2.5
3. Member has osteopenia with a high pre-treatment FRAX fracture probability (See Appendix B)

© 2019 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

C. Breast Cancer

Authorization of 12 months may be granted to members who are receiving adjuvant aromatase inhibitor therapy for breast cancer.

D. Prostate Cancer

Authorization of 12 months may be granted to members who are receiving androgen deprivation therapy for prostate cancer.

E. Glucocorticoid-induced Osteoporosis

Authorization of 12 months may be granted to members with glucocorticoid-induced osteoporosis when ALL of the following criteria are met:

1. Member is currently receiving or will be initiating glucocorticoid therapy
2. Member has had an oral bisphosphonate trial of at least 1-year duration OR there is a clinical reason to avoid treatment with an oral bisphosphonate (See Appendix A)
3. Member meets ANY of the following criteria:
 - a. Member has a history of a fragility fracture
 - b. Member has a pre-treatment T-score of ≤ -2.5
 - c. Member has osteopenia with a high pre-treatment FRAX fracture probability (See Appendix B)

III. CONTINUATION OF THERAPY

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

IV. APPENDIX

Appendix A. Clinical reasons to avoid oral bisphosphonate therapy

- Esophageal abnormality that delays emptying such as stricture of achalasia
- Active upper gastrointestinal problem (e.g., dysphagia, gastritis, duodenitis, erosive esophagitis, ulcers)
- Inability to stand or sit upright for at least 30 to 60 minutes
- Inability to take at least 30 to 60 minutes before first food, drink, or medication of the day
- Renal insufficiency (creatinine clearance <35 mL/min)
- History of intolerance to an oral bisphosphonate

Appendix B. WHO Fracture Risk Assessment Tool

- High FRAX fracture probability: 10 year major osteoporotic fracture risk $\geq 20\%$ or hip fracture risk $\geq 3\%$.
- 10-year probability; calculation tool available at: <http://www.shef.ac.uk/FRAX/tool.jsp>

V. REFERENCES

1. Prolia [package insert]. Thousand Oaks, CA: Amgen Inc.; May 2018.
2. The NCCN Drugs & Biologics Compendium™ © 2016 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed October 18, 2017.
3. Bisphosphonates. *Drug Facts and Comparisons. Facts & Comparisons® eAnswers* [online]. 2015. Available from Wolters Kluwer Health, Inc. Accessed October 18, 2017.
4. Cosman F, de Beur SJ, LeBoff MS, et al. National Osteoporosis Foundation. Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int*. 2014;25(10): 2359-2381.
5. Jeremiah MP, Unwin BK, Greenwald MH, et al. Diagnosis and management of osteoporosis. *Am Fam Physician*. 2015;92(4):261-268.
6. Watts NB, Bilezikian JP, Camacho PM, et al. American Association of Clinical Endocrinologists medical guidelines for clinical practice for the diagnosis and treatment of postmenopausal osteoporosis. *Endocr Pract*. 2016;22 (Suppl 4):1-42.

© 2019 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

7. ACOG Practice Bulletin Number 129: Osteoporosis. *Obstet Gynecol.* 2012;120(3):718-734.
8. National Institute for Health and Care Excellence. Osteoporosis Overview. Last updated August 2017. Available at: <http://pathways.nice.org.uk/pathways/osteoporosis>. Accessed October 18, 2017.
9. Treatment to prevent osteoporotic fractures: an update. Department of Health and Human Services, Agency for Healthcare Research and Quality. 2012; Publication No. 12-EHC023-EF. Available at www.effectivehealthcare.ahrq.gov/lbd.cfm.
10. Watts NB, Adler RA, Bilezikian JP, et al. Osteoporosis in men : an Endocrine Society clinical practice guideline. *J Clin Endocr Metab.* 2012;97(6):1802-1822.
11. Gralow JR, Biermann S, Farooki A, et al. NCCN Task Force Report: Bone Health in Cancer Care. *JNCCN.* 2013; 11(Suppl 3):S1-50.
12. FRAX[®] WHO fracture risk assessment tool. © World Health Organization Collaborating Centre for Metabolic Bone Diseases: University of Sheffield, UK. Available at: <http://www.shef.ac.uk/FRAX>. Accessed October 18, 2017.
13. Fink HA, Gordon G, Buckley L, et al. 2017 American College of Rheumatology Guidelines for the Prevention and Treatment of Glucocorticoid-Induced Osteoporosis. *Arthritis Care Res.* 2017;69:1521-1537.

© 2019 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.